

Algorithm for dealing with suspected SARS-CoV-2 coronavirus infections

If you are a person who has had close contact with someone who has been diagnosed with SARS-CoV-2 coronavirus infection (contact with a confirmed or probable case) or has been in a region where SARS-CoV-2 coronavirus transmission persists and you have experienced at least one of the symptoms of an acute respiratory infection listed:

- Fever
- Cough
- Souliness

YES

Call the SGGW clinic between 8:00 and 18:00. Don't come to the clinic!

If you have severe symptoms such as a temperature above 38 C, coughing, shortness of breath or breathing difficulties, you will most likely be referred to an infectious hospital or other specialist medical centre. The method of transport will be determined by your doctor in accordance with current health requirements. In the case of an emergency and serious health threat, call 112, 999 people directly to the infectious diseases ward (avoiding public transport and people's groups).

If you have small, common cold symptoms you will most likely be referred for treatment at home and asked to reduce your socio-professional activity for a period of several days after the symptoms have subsided (just stay at home). If nothing happens you can return to your daily activities on a date agreed with your doctor. If your condition worsens, you will have severe symptoms - temp. above 38 C, coughing, dyspnea or breathing difficulties - contact NZOZ SGGW by phone (do not come to the clinic!!!) or go directly to the infectious diseases ward (avoiding public transport and people's groups). In case of an emergency and serious health threat, call 112; 999

NO

Call the SGGW clinic between 8:00 and 18:00. Don't come to the clinic!

If you have no symptoms, you will most likely be asked to reduce your socio-occupational activity for a period of 14 days from exposure (just stay at home) and to keep yourself self-monitoring and measuring morning and evening temperatures during this period. If nothing happens during this period you can return to your daily activities. If your condition worsens, you will experience severe symptoms - temp above 38 C, coughing, shortness of breath or breathing difficulties